

Observational evidence of sensitivity of surface climate changes to land types and urbanization

Young-Kwon Lim¹, Ming Cai¹, Eugenia Kalnay², and Liming Zhou³

Department of Meteorology, the Florida State University

Department of Meteorology, University of Maryland

School of Earth and Atmospheric Sciences, Georgia Institute of Technology

Abstract. Sensitivity of surface climate change to land types is investigated for the Northern Hemisphere by subtracting the reanalysis from the observed surface temperature (observation minus reanalysis). The basis of this approach is that while reanalysis represents the large-scale climate changes due to greenhouse gases and atmospheric circulation, it is less sensitive to regional surface processes associated with land types.

OMR trends derived from two independent reanalyses (ERA40 and NNR) and two observation data sets (CRU and GHCN) show similar dependence upon land types, suggesting the attribution of OMRs to different land types is robust. OMR trends reveal 1) Warming over barren areas is larger than most other land types. 2) Urban areas show a large warming second only to barren areas. 3) Croplands with agricultural activity show a larger warming than natural broadleaf forests. The overall assessment indicates surface warming is larger for areas that are barren, anthropogenically developed, or covered with needle-leaf forests.

1. Introduction

Greenhouse gases [IPCC, 2001] and land-uses [Pielke *et al.*, 2002] are known as primary human impacts on climate change. They both contribute to surface warming and tend to reduce the diurnal temperature range [Gallo and Owen, 1999; Kalnay and Cai, 2003, hereafter KC; Kalnay *et al.* 2005]. It has been suggested that land-use impact on surface climate change is not negligible compared to atmospheric greenhouse effect [Lofgren, 1995; Bounoua *et al.*, 1999; KC].

However, an assessment of the local response to the land-use impact has not been fully addressed because there is no available way to reliably separate the local climate change signal from the global one. Only the urban impact has been estimated by comparing observations in cities with those in rural areas. The estimate, however, is limited to urban areas and provides little information about the impact of other land-cover types on the long-term surface temperature trend. In addition, estimates vary with the method of classifying urban and rural areas. For instance, the warming trend due to urbanization over the US estimated by Easterling *et al.* [1996] based on population is +0.06°C/century whereas Hansen *et al.* [2001] based on night-light observation obtain +0.15°C/century, with regional urban warming and cooling.

The objective of this study is to assess the surface warming sensitivity to land types and urbanization using the “Observation minus Reanalysis (OMR)” approach [KC; Zhou *et al.*, 2004; Frauenfeld *et al.*, 2005]. Specifically, we apply OMR to estimate the impact of urban areas and other land types for the Northern Hemisphere (NH) based on the comparison between trends observed in surface stations with those estimated with the surface temperature derived from the NCEP/NCAR reanalysis (NNR) [Kalnay *et al.*, 1996; Kistler *et al.*, 2001] and ECMWF-40 (ERA40) [Simmons *et al.*, 2004]. The OMR method relies on the fact that reanalyses represent the large-scale climate changes due to

greenhouse gases and atmospheric circulation, but the NNR and (to a lesser extent) the ERA40 are insensitive to regional surface processes associated with different land types [NRC, 2005]. Thus, the surface observations after removing the reanalysis enable us to isolate local near-surface warming patterns from the global-scale warming signal. We will attempt to attribute the local OMR surface warming patterns to land types using land type classifications made with satellite observations.

2. Data

The NH surface temperature data in this study consist of two gridded (2.5°×2.5°) reanalyses (ERA40 (2-meter temperature from <http://data.ecmwf.int/data/>) and NNR), and two gridded (5°×5°) observations (Global Historical Climatology Network (GHCN) [Peterson and Vose, 1997], and Climatic Research Unit (CRU) [Jones and Moberg, 2003]) from <http://www.ncdc.noaa.gov> and <http://www.cru.uea.ac.uk>, respectively. NCEP/DOE reanalysis version 2 (R2) is also used for investigating OMR time series in section 3. The rationale for the OMR approach is that the reanalysis is less sensitive to surface processes because little or no surface data or information about land-surface changes were used in the data assimilation process. The NNR creates its own estimate of surface fields from the upper air information combined with model parameterizations of surface processes. As a result, the NNR should not be sensitive to local surface properties at all, even if it should show climate change effects to the extent that they affect the observations above the surface [Kistler *et al.*, 2001]. Moreover, it has been shown that a reanalysis made with a frozen model (as the case of the NNR) can detect an anthropogenic trend present in observations assimilated by the reanalysis system essentially at its full strength [Cai and Kalnay, 2005]. As to the ERA40, surface temperature and soil moisture are estimated by assimilating the CRU observations in an off-line mode. Therefore, it is expected that the OMR using ERA40 would contain a portion of climate trend due to the impact of land types, resulting in a smaller OMR trend than that derived from NNR. However, since the surface air temperature observations are only used indirectly, it is expected that the OMR method applied to ERA40 also has some useful information about land types underneath.

To attribute OMR trend to land types, we used the Moderate Resolution Imaging Spectro-radiometer (MODIS) land cover classification map [Friedl *et al.*, 2002] from <http://edcdaac.usgs.gov/modis/mod12q1.asp>. The data consist of 16 land types with 1km×1km pixels.

3. Hemispheric surface temperature time series

Plotted in Fig. 1 are surface temperature anomalies averaged over the NH derived from three reanalyses and two observations. Anomalies are further adjusted to have zero mean over the last 10 years (1993-2002) because the biases of the reanalysis data for the most recent years are smallest [Simmons *et al.* 2004]. It should be

pointed out that the ERA40 time series in Fig. 1a is nearly identical to the top panel of Fig. 1 in Simmons et al. [2004]. It is seen that the two observational data sets (e.g. CRU and GHCN) are nearly indistinguishable (Figs 1a), showing a gradual warming trend over the NH. Reanalyses are in good agreement with the observations in terms of capturing the inter-annual variability and the long-term warming trends. As expected, the upward trend of the ERA40 is closer to observations than both the NNR and its follow-up, R2. Nevertheless, it is evident that the observations exhibit a larger warming trend compared to the reanalyses (Fig. 1a). As a result, OMRs show a positive trend (Fig. 1b), with a larger trend using NNR or R2 than ERA40. This suggests that OMR time series using ERA40, NNR and R2 support the NNR-based findings of KC, namely that the reanalysis trend is smaller than the observations' trend.




Fig. 1. Time series (three-year running mean) of (a) land surface temperature anomalies ($^{\circ}\text{C}$) derived from CRU, GHCN, ERA40, NNR, and R2 and (b) the OMRs. Anomaly values are obtained by removing the 30-yr mean from 1961 to 1990 and they are further adjusted to have zero mean over the last 10 years (1993-2002).

4. OMR trends with respect to land types

We now relate OMRs long-term trends to the surface properties. Areal fractions of individual land covers ($1\text{km}\times 1\text{km}$) are calculated for each $5^{\circ}\times 5^{\circ}$ grid, which is the same resolution as the surface temperature data. Displayed in Fig. 2a is the geographic distribution of the dominant land cover types, whose areal percentage in each grid exceeds at least 40% (equivalent to about $100,000\text{km}^2$ at 30°N). In order to avoid ambiguity in classifying dominant types we excluded grid boxes (colored black in Fig. 2a) where the dominant type covers less than 40% of the area. A more stringent requirement with higher areal percentage is not desirable because it would lead to a situation in which the number of the qualified grid points in each category is too small to draw any statistically significant results. Fig. 2a includes major land types characterizing the earth surface, as listed in Table 1. However, urban, wetland, closed shrub land, and natural vegetation mosaic are absent in Fig. 2a (and no color bar is assigned to these categories) because none of these 4 categories has the largest percentage coverage in any of the $5^{\circ}\times 5^{\circ}$ grid boxes. We will use the high-resolution MODIS data to assess the urban impact on the long-term surface temperature trend in the next section. Panel (b) displays the mean OMR trends and their vertical error bars at 95% significance level as a function of land type using GHCN/NNR (red) and GHCN/ERA40 (blue),

respectively. As in KC, the OMR trend per decade is obtained by taking the average of two decadal mean differences, that is, 90's - 80's and 70's - 60's, at each grid point, followed by averaging for the same land types. Table 1 lists the number of $5^{\circ}\times 5^{\circ}$ grid boxes used for the OMR trend calculation for each of the 16 land-cover categories. The OMR calculations were performed only in grid boxes where observation and reanalysis coexist.




Fig. 2. (a) Land cover map derived from MODIS. Grid boxes in which the dominant land cover type covers less than 40% are colored black and not used in the analysis presented in panel (b). (b) the mean OMR trend of “GHCN-minus-NNR” (red), and “GHCN-minus-ERA40” (blue) per decade ($^{\circ}\text{C}/\text{decade}$) over the NH as a function of land types. Filled squares represent the mean OMR trends and vertical lines the error bars at 95% significance level. The OMR trend per decade is obtained by taking the average of two decadal mean difference (90's - 80's and 70's - 60's).

Table 1. 16 Land-cover categories from MODIS and the number of $5^{\circ}\times 5^{\circ}$ grid boxes used for calculation of OMR trends per decade.

1) Evergreen needle-leaf forest	2) Evergreen broadleaf forest	3) Deciduous needle-leaf forest	4) Deciduous broadleaf forest
29	42	4	3
5) Mixed forest	6) Closed shrubland	7) Open shrubland	8) Woody Savannah
31	0	81	6
9) Savannah	10) Grassland	11) Wetland	12) Cropland
6	36	0	51
13) Urban	14) Natural vegetation mosaic	15) Snow and Ice	16) Barren or sparsely vegetated
0	0	3	56

The two independent reanalyses appear to reveal a very similar dependence of the OMR trends with respect to land types as well as their statistical significance levels (Fig. 2b). This suggests that the attribution of the OMR trends to different land types is robust. The key features in the OMR trends are summarized below:

(i) The OMR trend over barren areas (category 16) ($\geq 0.3^{\circ}\text{C}/\text{decade}$) is larger than most of the other land types. It is known that the evaporation feedback decreases the surface warming. The results seem to suggest that over barren or arid areas where soil moisture is very limited, the evaporation feedback would be negligible, explaining a larger local surface warming under the same amount of radiative forcings due to anthropogenic greenhouse gases, as discussed in Dai et al. [2004] and Hales et al. [2004].

(ii) The OMR over croplands and grass (12 and 10), where a large seasonal vegetation change takes place, show a moderate decadal warming ($\sim 0.2^\circ\text{C}/\text{decade}$). In contrast, for the land type 4 (broadleaf deciduous), which experiences a similarly large seasonal variation in terms of vegetation growth as the cropland, the OMR trend is very small. This suggests human activities could be responsible for an additional local warming.

(iii) In addition, the positive OMR trend is evident over arid areas with shrubs (7) ($\sim 0.2^\circ\text{C}/\text{decade}$). Together with a result in (i), this suggests that the OMR trend could be related to the soil moisture level [NRC, 2005].

(iv) There are large OMR trends over the needle-leaf forests (1 and 3) ($\geq 0.2^\circ\text{C}/\text{decade}$), and conversely, broadleaf tree areas (2 and 4) do not show a significant warming trend ($0.05^\circ\text{C}/\text{decade}$). These results are quite consistent with modeling works [Shukla *et al.*, 1990; Xue and Shukla, 1993; Giambelluca *et al.*, 1997] which show that reduced (increased) transpiration by clearing (creating) broadleaf forest causes warming (cooling). We therefore suggest that larger transpiration and evaporative cooling over broadleaf forests may be related to weaker warming than needle-leaf forests.

5. Urban impact




Fig. 3. a) Geographical distribution of urban grids ($5^\circ \times 5^\circ$). Grid boxes where the fractional area of $1\text{km} \times 1\text{km}$ urban pixels is greater than 0.043 (in red), and between 0.01 and 0.042 (in blue) are categorized as big (small) urban areas. Time series ($^\circ\text{C}$) of b) GHCN-NNR, and c) CRU-NNR, for the areas of big urban areas (red solid), small urban areas (red dashed), agriculture (blue solid), natural broadleaf (blue dashed), and barren areas (black solid), respectively.

Urban impact on surface climate change is also assessed, along with the comparison with impacts of other land types. Grid boxes in which the percentage area of $1\text{km} \times 1\text{km}$ urban pixels exceed 4.3% (between 1 and 4.2%) are categorized as “big” (“small”) urban areas, so that the well-known large cities have been included in the big urban categories (Fig. 3a). 4.3%-urban pixels in a grid box of $5^\circ \times 5^\circ$ to an area of approximately $11,000\text{km}^2$ at 30°N . Major grid boxes categorized as big urban areas based on this configuration include France, Germany, Italy, and UK in Europe; Beijing, Guangzhou-Hongkong, Japan, Shanghai, South Korea, and Taiwan

in east Asia; California, Chicago, Florida, New York-Boston in North America (Fig. 3a). We then examine whether OMR trends over these “extended urban areas” are indeed larger than other areas.

Figure 3 depicts the OMR time series averaged over big urban areas, small urban areas, agriculture, broadleaf forests, and barren areas. We can find that the order of magnitude in warming trends, shown in both panels is consistent. Big urban area shows a larger warming trend than small urban and agricultural areas. However, urban warming is weaker than that over barren areas, which are the most sensitive regions in terms of surface warming. OMR decadal warming trends (computed with the same method as in section 4) are 0.24, 0.19, 0.18, 0.03, 0.33 for these five categories using “GHCN-NNR”. The trends using “CRU-NNR” are 0.21, 0.19, 0.2, 0.02, 0.34, respectively. It appears that the barren or urban surface with limited soil moisture exerts a strong surface warming response due to a weaker evaporative cooling process [Hales *et al.*, 2004; NRC, 2005]. In contrast, warming trend over natural broadleaf forests is smaller than other land types, supporting the notion that the high soil moisture damps surface warming.

6. Summary and discussion

Prominent features identified from the OMR using reanalysis (NNR or ERA-40) and observational datasets (GHCN or CRU) are

1) Barren areas show a larger warming trend ($\geq 0.3^\circ\text{C}/\text{decade}$) than most of other land types. It should be pointed out that the larger warming over the barren areas may not necessarily be associated with land-cover change, such as desertification or deforestation. For example, large OMR difference over Sahara desert may simply imply that the local surface warming due to greenhouse effects are stronger there because of a much weaker evaporation feedback over the desert [Dai *et al.* 2004; Hales *et al.*, 2004]. Diffenbaugh [2005] and NRC [2005] discussed that greenhouse gas-induced atmosphere-land cover feedbacks can increase or decrease the local surface temperature by modifying surface albedo, the soil moisture and evaporation process.

2) Among all other land types, urban impact on surface warming is strongest second to the barren areas. In particular, it exceeds the warming in agricultural land by $0.06^\circ\text{C}/\text{decade}$.

3) Within areas of strong seasonality, agricultural areas (cropland) and mixed forests show a moderate warming trend ($\sim 0.2^\circ\text{C}/\text{decade}$) whereas natural deciduous broad-leaf areas do not. But this may need further reassessment because of the small number of samples for the deciduous broad-leaf class, even if its error bar is very small.

4) In contrast, for the highly vegetated areas such as tropical forest and low-latitude evergreen broad-leaf tree areas, the strong evaporative cooling appears to suppress the surface warming ($< 0.1^\circ\text{C}/\text{decade}$). Dai *et al.* [2004] and Kumar *et al.* [2004] address that surface temperature change is related to the soil moisture, which in turn associated with vegetation types. We suggest that this reasoning is also applicable to explain much larger warming trends in mid-latitude needle-leaf forests than moist broadleaf forests identified in this study.

The results shown here support our hypothesis that the OMR can be attributed to the sensitivity of surface climate change to land types. The findings are mostly insensitive to the choice of reanalysis (NNR or ERA-40) or observational data (GHCN or CRU). It is important to note that this study uses the land cover estimated by recent MODIS observations, which may or may not reflect changes in land types. Therefore, part of the sensitivity of

the OMR to land types may be due to changes in land type. There may be also other uncertainties in OMR differences, but the fact that the OMR does provide a statistically significant and physically explainable sensitivity to land type suggests these uncertainties may be random and cancel out after averaging over a large number of grids with same land type. However, it is important to use climate modeling techniques for further understanding of the warming sensitivity to land types and the climate impact of land use changes.

7. Acknowledgement

This study is supported by the National Science Foundation (ATM-0403211 and ATM-0403518) and by the Climate Change Data and Detection program of NOAA Office of Global Programs (GC04-259). ERA40 data were obtained from the ECMWF public web server, <http://data.ecmwf.int/data>. The constructive comments and suggestions from the anonymous reviewer are helpful.

References

- Bounoua, L., and Coauthors (1999), Interactions between vegetation and climate: Radiative and physiological effects of doubled atmospheric CO₂, *J. Clim.*, *12*, 309-323.
- Cai, M., and E. Kalnay (2005), Can reanalysis have anthropogenic climate trends without model forcing?, *J. Clim.*, in press.
- Dai, A, K. E. Trenberth, and T. Qian (2004), A global dataset of Palmer drought severity index for 1870-2002: Relationship with soil moisture and effects of surface warming, *J. Hydrometeorol.*, *5*, 1117-1130.
- Diffenbaugh, N. S. (2005), Atmosphere-land cover feedbacks alter the response of surface temperature to CO₂ forcing in the western United States, *Clim. Dyn.*, *24*, 237-251.
- Easterling, D. R., T. C. Peterson, and T. R. Karl (1996), On the development and use of homogenized climate datasets, *J. Clim.*, *9*, 2941-2944.
- Frauenfeld, O. W., T. Zhang, and M. C. Serreze (2005), Climate change and variability using European Center for Medium-Range Weather Forecast analysis (ERA-40) temperatures on the Tibetan Plateau, *J. Geophys. Res.*, *110*, D02101, doi:10.1029/2004JD005230.
- Friedl, M. A., and Coauthors (2002), Global land cover from MODIS: Algorithms and early results, *Remote Sens. Environ.*, *83*, 287-302.
- Gallo, K. P., and T. W. Owen (1999), Satellite-based adjustments for urban heat island temperature bias, *J. Appl. Meteorol.*, *38*, 806-813.
- Giambelluca, T. W., D. Hölscher, T. X. Bastos, R. R. Frazão, M. A. Nullet, and A. D. Ziegler (1997), Observations of albedo and radiation balance over postforest land surfaces in the eastern Amazon basin, *J. Clim.*, *10*, 919-928.
- Hales, K., J. D. Neelin, and N. Zeng (2004), Sensitivity of tropical land climate to leaf area index: Role of surface conductance versus albedo, *J. Clim.*, *17*, 1459-1473.
- Hansen, J. E., R. Ruedy, M. Sato, M. Imhoff, W. Lawrence, D. Easterling, T. Peterson, and T. Karl (2001), A closer look at United States and global surface temperature change, *J. Geophys. Res.*, *106*, 23947-23963.
- IPCC (2001), *Climate Change 2001: The scientific basis*, Cambridge University Press.
- Jones, P. D., and A. Moberg (2003), Hemispheric and large-scale surface air temperature variations: An extensive revision and an update to 2001, *J. Clim.*, *16*, 206-223.
- Kalnay, E., and co-authors (1996), The NCEP/NCAR 40-year reanalysis project, *Bull. Amer. Meteor. Soc.*, *77*, 437-471.
- Kalnay, E., and M. Cai (2003), Impact of urbanization and land-use on climate change, *Nature*, *423*, 528-531.
- Kalnay, E., M. Cai, H. Li, and C. J. Tobin (2005): Estimation of the impact of land-surface forcings on temperature trends in eastern United States. *J. Geophys. Res.* (accepted).
- Kistler, R., and Coauthors (2001), The NCEP/NCAR 50-year reanalysis: monthly means CD-ROM and documentation, *Bull. Amer. Meteor. Soc.*, *82*, 247-267.
- Kumar, A., F. Yang, L. Goddard, and S. Schubert (2004), Differing trends in the tropical surface temperatures and precipitation over land and oceans, *J. Clim.*, *17*, 653-664.
- Lofgren, B. M. (1995), Sensitivity of land-ocean circulations, precipitation, and soil moisture to perturbed land surface albedo. *J. Clim.*, *8*, 2521-2542.
- National Research Council (2005), *Radiative forcing of climate change: Expanding the concept and addressing uncertainties*, 207pp., National Academies Press.
- Peterson, T. C., and R. S. Vose (1997), An overview of the Global Historical Climatology Network temperature database, *Bull. Amer. Meteor. Soc.*, *78*, 2837-2848.
- Pielke Sr, R. A., and Coauthors (2002), The influence of land-use change and landscape dynamics on the climate system: relevance to climate-change policy beyond the radiative effects of greenhouse gases, *Phil. Trans. Roy. Soc. Lond.*, *360*, 1-5.
- Shukla, J., C. Nobre, and P. Sellers (1990), Amazon deforestation and climate change, *Science*, *247*, 1322-1325.
- Simmons, A. J., P. D. Jones, V. da Costa Bechtold, A. C. M. Beljaars, P. W. Kallberg, S. Saarinen, S. M. Uppala, P. Viterbo, and N. Wedi (2004), Comparison of trends and low-frequency variability in CRU, ERA-40, and NCEP/NCAR analyses of surface air temperature, *J. Geophys. Res.*, *109*, D24115, doi:10.1029/2004JD005306.
- Xue, Y., and J. Shukla (1993), The influence of land surface properties on Sahel climate. Part I: desertification, *J. Clim.*, *6*, 2232-2245.
- Zhou, L., R. E. Dickinson, Y. Tian, J. Fang, Q. Li, R. K. Kaufmann, C. J. Tucker, and R. B. Myneni (2004), Evidence for a significant urbanization effect on climate in China, *Proceedings of the National Academy of Sciences*, *101*, 9540-9544.